

On the number of hamiltonian cycles in triangulations with few separating triangles

Gunnar Brinkmann Annelies Cuvelier Jasper Souffriau
Nico Van Cleemput

Combinatorial Algorithms and Algorithmic Graph Theory
Department of Applied Mathematics, Computer Science and Statistics
Ghent University




Outline

- 1 Introduction
 - Definitions
 - Known results
- 2 Technique
 - Counting base
 - Subgraphs
 - Partitions
 - One separating triangle
- 3 Results
 - New bounds
 - Conjectured bounds
 - Summary




Plane triangulation

A (plane) triangulation is a plane graph in which each face is a triangle.



Hamiltonian cycle

A hamiltonian cycle C in a graph $G = (V, E)$ is a spanning subgraph of G which is isomorphic to a cycle.




Hamiltonian cycle

\mathcal{H}_G : set of all hamiltonian cycles in graph G




Separating triangle

A separating triangle S in a triangulation G is a subgraph of G which is isomorphic to C_3 such that $G - S$ has two components.




Separating triangle

A separating triangle S in a triangulation G is a subgraph of G which is isomorphic to C_3 such $G - S$ has two components.



4-connected triangulation

A triangulation is 4-connected if and only if it contains no separating triangles.



Lower bound on number of hamiltonian cycles

Theorem (Whitney, 1931)

Every 4-connected triangulation is hamiltonian (i.e., contains at least one hamiltonian cycle).

Theorem (Jackson and Yu, 2002 (reformulated))

Every triangulation with at most 3 separating triangles is hamiltonian (i.e., contains at least one hamiltonian cycle).



Lower bound on number of hamiltonian cycles

Theorem (Kratochvíl and Zeps, 1988)

Every hamiltonian triangulation on at least 5 vertices contains at least four hamiltonian cycles.

Theorem (Hakimi, Schmeichel and Thomassen, 1979)


Every 4-connected triangulation on n vertices contains at least $\frac{n}{\log_2 n}$ hamiltonian cycles.



Lower bound on number of hamiltonian cycles

Conjecture (Hakimi, Schmeichel and Thomassen, 1979)

Every 4-connected triangulation on n vertices contains at least $2(n-2)(n-4)$ hamiltonian cycles.



Lower bound on number of hamiltonian cycles

Theorem (Hakimi, Schmeichel and Thomassen, 1979)


Every 4-connected triangulation on n vertices contains at least $\frac{n}{\log_2 n}$ hamiltonian cycles.




Proof by Hakimi, Schmeichel and Thomassen



Proof by Hakimi, Schmeichel and Thomassen



Proof by Hakimi, Schmeichel and Thomassen



Zigzag

Proof by Hakimi, Schmeichel and Thomassen

For each edge v, w in G : pick hamiltonian cycle containing $xvwy$.

$\Rightarrow \leq 3n - 6$ hamiltonian cycles.


Each hamiltonian cycle occurs at most α times.

$$\Rightarrow |\mathcal{H}_G| \geq \frac{3n-6}{\alpha}$$



Proof by Hakimi, Schmeichel and Thomassen

Let H be hamiltonian cycle that occurs α times.




At least $\frac{\alpha}{3}$ zigzags intersect in at most one vertex.



Proof by Hakimi, Schmeichel and Thomassen

New hamiltonian cycle for each independent zigzag switch.



$$\Rightarrow |\mathcal{H}_G| \geq 2^{\frac{\alpha}{3}}$$

Proof by Hakimi, Schmeichel and Thomassen

$$\log_2 |\mathcal{H}_G| \geq \frac{\alpha}{3} \geq \frac{n-2}{|\mathcal{H}_G|}$$

↓

$$|\mathcal{H}_G| \log_2 |\mathcal{H}_G| \geq n - 2$$

↓

$$|\mathcal{H}_G| > \frac{n}{\log_2 n}$$



Outline

- 1 Introduction
 - Definitions
 - Known results
- 2 **Technique**
 - Counting base
 - Subgraphs
 - Partitions
 - One separating triangle
- 3 Results
 - New bounds
 - Conjectured bounds
 - Summary







Counting base (S, r) for \mathcal{H} in G

General technique for finding a lower bound for the size of an arbitrary set \mathcal{H} of hamiltonian cycles in a given graph G .




Counting base (S, r) for \mathcal{H} in G

- $S \subseteq \{\text{subgraphs of } G\}$






Counting base (S, r) for \mathcal{H} in G

- $S \subseteq \{\text{subgraphs of } G\}$
- $r : S \rightarrow \{\text{subgraphs of } G\}$




Counting base (S, r) for \mathcal{H} in G

- $S \subseteq \{\text{subgraphs of } G\}$
- $r : S \rightarrow \{\text{subgraphs of } G\}$
- $\mathcal{H} \subseteq \mathcal{H}_G$




Counting base (S, r) for \mathcal{H} in G

- 1 Each $s \in S$ must be contained in at least one $C \in \mathcal{H}$.




Counting base (S, r) for \mathcal{H} in G

- 1 Each $s \in S$ must be contained in at least one $C \in \mathcal{H}$.
- 2 For each $s \in S$ we have that $s \not\subseteq r(s)$.




Counting base (S, r) for \mathcal{H} in G

- 1 Each $s \in S$ must be contained in at least one $C \in \mathcal{H}$.
- 2 For each $s \in S$ we have that $s \not\subseteq r(s)$.
- 3 For each $s \in S$ and $C \in \mathcal{H}$ with $s \subseteq C$ we have that $(C \setminus s) \cup r(s) \in \mathcal{H}$.



C





$(C \setminus s) \cup r(s)$





Counting base (S, r) for \mathcal{H} in G

- 1 Each $s \in S$ must be contained in at least one $C \in \mathcal{H}$.
- 2 For each $s \in S$ we have that $s \not\subseteq r(s)$.
- 3 For each $s \in S$ and $C \in \mathcal{H}$ with $s \subseteq C$ we have that $(C \setminus s) \cup r(s) \in \mathcal{H}$.
- 4 For two different $s_1, s_2 \in S$ and for any $C \in \mathcal{H}$ containing both subgraphs we have that $(C \setminus s_1) \cup r(s_1) \neq (C \setminus s_2) \cup r(s_2)$.


 $s_1 \subseteq C$

 $s_2 \subseteq C$

Counting base (S, r) for \mathcal{H} in G

- 1 Each $s \in S$ must be contained in at least one $C \in \mathcal{H}$.
- 2 For each $s \in S$ we have that $s \not\subseteq r(s)$.
- 3 For each $s \in S$ and $C \in \mathcal{H}$ with $s \subseteq C$ we have that $(C \setminus s) \cup r(s) \in \mathcal{H}$.
- 4 For two different $s_1, s_2 \in S$ and for any $C \in \mathcal{H}$ containing both subgraphs we have that $(C \setminus s_1) \cup r(s_1) \neq (C \setminus s_2) \cup r(s_2)$.


 $(C \setminus s_1) \cup r(s_1)$

 $(C \setminus s_2) \cup r(s_2)$

Overlap $o_S(C, x)$ and maximum overlap $O_S(\mathcal{H}, r)$


- $\bar{S} = S \cup \{r(s) \mid s \in S\}$




Overlap $o_S(C, x)$ and maximum overlap $O_S(\mathcal{H}, r)$

- $\bar{S} = S \cup \{r(s) \mid s \in S\}$
- For each $x \in \bar{S}$ and each $C \in \mathcal{H}$ with $x \subseteq C$:

$$o_S(C, x) = |\{s \in S \mid x \cap s \neq \emptyset \text{ and } s \subseteq C\}|$$



$x \subseteq C$




$s \subseteq C$


Overlap $o_S(C, x)$ and maximum overlap $O_S(\mathcal{H}, r)$

- $\bar{S} = S \cup \{r(s) \mid s \in S\}$
- For each $x \in \bar{S}$ and each $C \in \mathcal{H}$ with $x \subseteq C$:

$$o_S(C, x) = |\{s \in S \mid x \cap s \neq \emptyset \text{ and } s \subseteq C\}|$$



$x \subseteq C$



$s \subseteq C$

- $O_S(\mathcal{H}, r) = \max\{o_S(C, x) \mid C \in \mathcal{H}, x \in \bar{S} : x \subseteq C\}$

Theorem (Brinkmann, Souffriau, NVC, 2014)

Given a graph G , a set $\mathcal{H} \subseteq \mathcal{H}_G$, and a nonempty counting base (S, r) for \mathcal{H} , then

$$|\mathcal{H}| \geq \frac{|S|}{O_S(\mathcal{H}, r)}.$$





Zigzags

- Let G be a 4-connected triangulation
- Let S be the set of all zigzags ($|S| = 6n - 12$)
- r switches zigzag to mirror image ($\bar{S} = S$)
- $\mathcal{H} = \mathcal{H}_G$
- $o_S(C, x) \leq 5$, so $O_S(\mathcal{H}_G, r) \leq 5$

$$\Rightarrow |\mathcal{H}_G| \geq \frac{6n-12}{O_S(\mathcal{H}_G, r)} \geq \frac{6n-12}{5}$$



Root path and inverse root path



Theorem (Brinkmann, Souffriau, NVC, 2014)

Every 4-connected triangulation on n vertices has $12(n - 2)$ root paths and inverse root paths.

Root path and inverse root path


- Let G be a 4-connected triangulation
- Let S be the set of all root paths and inverse root paths ($|S| = 12(n - 2)$)
- r switches root path to inverse root path on same vertices (and vice versa) ($\bar{S} = S$)
- $\mathcal{H} = \mathcal{H}_G$

This gives a counting base (S, r) for \mathcal{H}_G .




Theorem (Brinkmann, Souffriau, NVC, 2014)

For each 4-connected triangulation G we have that $O_S(\mathcal{H}_G, r) \leq 5$ with S the set of all root and inverse root paths in G .




Theorem (Brinkmann, Souffriau, NVC, 2014)

For each 4-connected triangulation G we have that $O_S(\mathcal{H}_G, r) \leq 5$ with S the set of all root and inverse root paths in G .




Theorem (Brinkmann, Souffriau, NVC, 2014)

For each 4-connected triangulation G we have that $O_S(\mathcal{H}_G, r) \leq 5$ with S the set of all root and inverse root paths in G .




Theorem (Brinkmann, Souffriau, NVC, 2014)

For each 4-connected triangulation G we have that $O_S(\mathcal{H}_G, r) \leq 5$ with S the set of all root and inverse root paths in G .



Theorem (Brinkmann, Souffriau, NVC, 2014)

For each 4-connected triangulation G we have that $O_S(\mathcal{H}_G, r) \leq 5$ with S the set of all root and inverse root paths in G .



Theorem (Brinkmann, Souffriau, NVC, 2014)

Every 4-connected triangulation on n vertices has at least $\frac{12}{5}(n - 2)$ hamiltonian cycles.



Maximum overlap of a subset S_i of S

Let (S, r) be a counting base for \mathcal{H} , and let S_i be a subset of S :

$$O_{S|S_i}(\mathcal{H}, r) = \max\{o_S(C, s) \mid C \in \mathcal{H}, s \in S_i : s \subseteq C\}$$



Theorem (Brinkmann, Cuvelier, NVC, 2015)

Given a graph G , a set $\mathcal{H} \subseteq \mathcal{H}_G$, a nonempty counting base (S, r) for \mathcal{H} , and a partition S_1, S_2, \dots, S_k of S , then

$$|\mathcal{H}| \geq \sum_{i=1}^k |S_i| \frac{2}{O_S(\mathcal{H}, r) + O_{S_i}(\mathcal{H}, r)}.$$




Theorem (Brinkmann, Cuvelier, NVC, 2015)


Every 4-connected triangulation on n vertices has at least $\frac{161}{60}(n - 2)$ hamiltonian cycles.




Hour glasses



Hour glasses



Inverse hour glass



Sparse set of hour glasses for a triangle t

Definition

A set H of hour glasses is a sparse set for a triangle t if

- no two elements $h \neq h'$ have the same set of v-edges, and
- for each $h \in H$: if t is not one of the two triangles of h , then h contains no edge of t as v-edge or base edge.

Lemma

A 4-connected triangulation with n vertices contains a sparse set of hour glasses of size $6n - 21$ for each facial triangle.



Set of hamiltonian cycles sharing an edge with a face

\mathcal{H}_G^t : set of all hamiltonian cycles in graph G that share an edge with the face t .

- Let G be a 4-connected triangulation
- Let t be a face of G
- Let S be a sparse set of hour glasses for t
- r maps an hour glass to its inverse

This gives a counting base (S, r) for \mathcal{H}_G^t .



Set of hamiltonian cycles sharing an edge with a face

The overlap for hour glasses is at most 4, so...

$$\Rightarrow |\mathcal{H}_G^t| \geq \frac{6n-21}{O_S(\mathcal{H}_G^t, r)} \geq \frac{6n-21}{4}$$






One separating triangle

Theorem (Brinkmann, Souffriau, NVC, 2014)

Every 3-connected triangulation on n vertices with exactly one separating triangle has at least $\frac{6n-27}{4}$ hamiltonian cycles.





 T_0  T_i

With T_0 at least as many vertices as T_i .

- Both T_o and T_i are K_4 .




- Both T_o and T_i are K_4 .
- T_i is K_4




- Both T_o and T_i are K_4 .
- T_i is K_4
 - T_o is 4-connected and has $n - 1$ vertices
 - T_o has at least $\frac{6(n-1)-21}{4} = \frac{6n-27}{4}$ hamiltonian cycles sharing an edge with the separating triangle



- Both T_o and T_i are K_4 .
- T_i is K_4
 - T_o is 4-connected and has $n - 1$ vertices
 - T_o has at least $\frac{6(n-1)-21}{4} = \frac{6n-27}{4}$ hamiltonian cycles sharing an edge with the separating triangle



- Both T_o and T_i are K_4 .
- T_i is K_4
 - T_o is 4-connected and has $n - 1$ vertices
 - T_o has at least $\frac{6(n-1)-21}{4} = \frac{6n-27}{4}$ hamiltonian cycles sharing an edge with the separating triangle




- Both T_o and T_i are K_4 .
- T_i is K_4
- T_i is 4-connected




- Both T_o and T_i are K_4 .
- T_i is K_4
- T_i is 4-connected
 - T_o is 4-connected and has at least $\frac{n+3}{2}$ vertices
 - T_o has at least $\frac{6\frac{n+3}{2}-21}{4} = \frac{3n-12}{4}$ hamiltonian cycles sharing an edge with the separating triangle




- Both T_o and T_i are K_4 .
- T_i is K_4
- T_i is 4-connected
 - T_o is 4-connected and has at least $\frac{n+3}{2}$ vertices
 - T_o has at least $\frac{6\frac{n+3}{2}-21}{4} = \frac{3n-12}{4}$ hamiltonian cycles sharing an edge with the separating triangle



- Both T_o and T_i are K_4 .
- T_i is K_4
- T_i is 4-connected
 - T_o is 4-connected and has at least $\frac{n+3}{2}$ vertices
 - T_o has at least $\frac{6\frac{n+3}{2}-21}{4} = \frac{3n-12}{4}$ hamiltonian cycles sharing an edge with the separating triangle



- Both T_o and T_i are K_4 .
- T_i is K_4
- T_i is 4-connected
 - T_o is 4-connected and has at least $\frac{n+3}{2}$ vertices
 - T_o has at least $\frac{6\frac{n+3}{2}-21}{4} = \frac{3n-12}{4}$ hamiltonian cycles sharing an edge with the separating triangle



- Both T_o and T_i are K_4 .
- T_i is K_4
- T_i is 4-connected
 - T_o is 4-connected and has at least $\frac{n+3}{2}$ vertices
 - T_o has at least $\frac{6\frac{n+3}{2}-21}{4} = \frac{3n-12}{4}$ hamiltonian cycles sharing an edge with the separating triangle
 - original triangulation has at least $2\frac{3n-12}{4} = \frac{6n-24}{4} > \frac{6n-27}{4}$ hamiltonian cycles

Outline

- 1 Introduction
 - Definitions
 - Known results
- 2 Technique
 - Counting base
 - Subgraphs
 - Partitions
 - One separating triangle
- 3 Results
 - New bounds
 - Conjectured bounds
 - Summary



Theorem (Brinkmann, Cuvelier, NVC, 2015)

Every 4-connected triangulation on n vertices has at least $\frac{161}{60}(n - 2)$ hamiltonian cycles.

Theorem (Brinkmann, Souffriau, NVC, 2014)

Every 3-connected triangulation on n vertices with exactly one separating triangle has at least $\frac{6n-27}{4}$ hamiltonian cycles.



Computational results

Conjectured bound for 4-connected triangulations by Hakimi, Schmeichel and Thomassen verified **up to 25** vertices.



Computational results

For 6 separating triangles it is known that there exist non-hamiltonian 3-connected triangulations.

Minimum number of hamiltonian cycles for 3-connected triangulations with at most 5 separating triangles computed **up to 23** vertices.



Computational results


	1	2	3	4	5
5	6	-	-	-	-
6	-	10	-	-	-
7	24	-	12	-	-
8	42	26	-	6	-
9	64	36	24	-	8
10	90	46	33	12	-
11	120	56	41	14	12
12	154	66	49	14	10
13	192	76	57	14	10
14	234	86	65	14	10
15	280	96	73	14	10
16	330	106	81	14	10
17	384	116	89	14	10
18	442	126	97	14	10
19	504	136	105	14	10
20	570	146	113	14	10
21	640	156	121	14	10
22	714	166	129	14	10
23	792	176	137	14	10

Computational results








	1	2	3	4	5
For $n \geq 12$	$2(n-1)(n-5)$	$10n-54$	$8n-47$	14	10



Extremal graphs



Extremal graphs



Summary

Lower bounds for the number of hamiltonian cycles in triangulations with few separating triangles

# sep. triangle	Old bound	New bound	Conjectured bound
0	$\frac{n}{\log_2 n}$	$\frac{161}{60}(n-2)$	$2(n-2)(n-4)$
1	4	$\frac{6n-27}{4}$	$2(n-1)(n-5)$
2	4	$[4, 10n-54]$	$10n-54$
3	4	$[4, 8n-47]$	$8n-47$
4	0	$[0, 14]$	14
5	0	$[0, 10]$	10